

## A NEW SPECIES OF *RASBORA* BLEEKER (CYPRINIFORMES: CYPRINIDAE) FROM MANIPUR, INDIA<sup>1</sup>

WAIKHOM VISHWANATH<sup>2,3</sup> AND JULIANA LAISRAM<sup>2,4</sup>

<sup>1</sup> Accepted July, 2002

<sup>2</sup> Department of Life Sciences, Manipur University, Canchipur 795 003, Manipur, India. )

<sup>3</sup> Email: vnath\_w@hotmail.com

<sup>4</sup> Email: pinkylaisram@yahoo.com

A new fish species of the genus *Rasbora* Bleeker is described from the Lokchao and Chatrickong rivers (Chindwin drainage) of Manipur, India. It is distinguished from *Rasbora rasbora* (Hamilton-Buchanan), the only other known species from the region, by the lateral line incomplete vs. complete, lateral transverse scale 4Y:Z/1/2 vs. 4Y:z/III, greater head length of 29.3(27.5-31.1) vs. 24.5(22.1-26.9)% of SL, greater gape width of 8.1 (7.1-9.1) vs. 6.8(6.0-7.6)% of SL and gill rakers numbering 4-5+ 11 vs. 3+9-10.

**Key words:** New fish species, *Rasbora*, Manipur

### INTRODUCTION

Freshwater fishes of the genus *Rasbora* Bleeker belong to subfamily Danioninae (=Rasborinae). Certain rasboras are much prized aquarium fishes (Brittan 1954). Four species of the genus are known from India, namely *R. caverii* (Jerdon), *R. daniconius* (Hamilton-Buchanan), *R. labiosa* Mukerji and *R. rasbora* (Hamilton-Buchanan). Hora (1921) reported *R. rasbora* (Hamilton-Buchanan) from Dhaneshwori stream near Dimapur, Nagaland, which forms part of the Brahmaputra drainage system. Further, Hora and Mukerji (1935) reported the species to be widely distributed. Vishwanath *et al.* (1998) listed *R. rasbora* from Chatrickong river, Ukhrul district, Manipur. A fish collection from the Lokchao river (a tributary of the Yu River of Myanmar which, in turn, flows into the Chindwin river) in Chandel district of Manipur and from Chatrickong river (a tributary of Chindwin river) in Ukhrul district, included 14 specimens of *Rasbora* which did not fit into any of the hitherto described species. The species is described here as a new species.

### MATERIAL AND METHODS

Measurements and counts followed Brittan (1954). Measurements were made with dial callipers to the nearest 0.1 mm and expressed in percentages of standard length (SL) and head length (HL). Type specimens are deposited in the Manipur University Museum of Fishes (MUMF).

*Rasbora ornatus* sp. novo (Fig. 1)

Holotype: MUMF 3032, 56.0 mm SL, Lokchao R., Moreh, Manipur, a tributary of the Yu River, (Chindwin drainage), 24.iii. 1999, W. Vishwanath Singh and party.

**Paratypes:** MUMF: 1210-1212, 3 specimens, 44.2-57.1 mm SL, Chatrickong R., 6.vi.1996, K. Selim; MUMF 3033/ 10, 10 specimens, 35.1-77.6 mm SL, same data as holotype.

**Diagnosis:** A medium-sized *Rasbora* with an incomplete lateral line. Lateral transverse scales 4½/1/2. Lips simple, first dorsal fin ray without a fleshy sheath and the lateral stripe on body running from tip of snout to the end of median caudal rays.

### DESCRIPTION

D. ii, 7; P. i, 12-13; V. i, 8; A. iii, 5; C. 9+8. Body elongate, slightly compressed, its depth 28.0 (21.1-29.3)% SL. Mouth small, cleft oblique, lip simple, lower jaw slightly prominent with an upward projecting knob at symphysis fitting into a corresponding depression in upper jaw. Barbels absent, pharyngeal teeth in three rows 2,4,5-5,4,2. Gill rakers soft and low, 4-5+ II. Branchiostegal rays 3. Lateral line with a downward curvature anteriorly, runs along below centre of body and terminates at a line vertical to anterior origin, or sometimes at posterior extremity of dorsal fin. Lateral line incomplete, 26-28 scales in longitudinal series with 11-20 scales perforated. Lateral transverse scales from dorsal fin origin to the lateral line 4Y2 and from lateral line to origin of pelvic 2. Dorsal fin inserted a little behind origin of pelvic fin, nearer to




Fig. 1: Lateral view of *Rasbora ornatus* sp. novo

NEW DESCRIPTIONS

caudal fin base than to tip of snout. Pelvic fin not reaching base of anal fin. Head moderate, eyes large, visible from ventral side. Inter-orbital space wide, slightly arched, 35.7(29.8-39.0)% HL. Inter-narial space 24.6(20.8-27.6)% HL. Mouth gape wide. Dorsal-hypural distance when carried forward, falling at anterior rim of eye. Sexual dimorphism unknown.

**Colour:** Body creamish, abdomen pale, dorsal part of head brownish. A deep dark blue longitudinal stripe extending from tip of snout to end of median caudal rays. Scales on lateral and dorsal sides of body with a dark edge formed by a row of spots on each scale. All fins dusky in appearance.

**Etymology:** The species is named after its beautiful colouration.

**Distribution:** INDIA: Manipur, Lokchao R., Moreh, Chatrirkong R.

**Remarks:** *Rasbora ornatus* sp. novo belongs to the Daniconius-complex as it has 14 circumpeduncular scales, 2 scales between lateral line and pelvic fin origin, the typical characters of the complex. As in some forms of this complex, this species shows reduced number of lateral line pores and prominent dark lateral stripe, which runs from the tip of the snout to the median caudal rays.

*Rasbora ornatus* sp. novo differs from *R. rasbora* in having an incomplete lateral line vs. complete and lateral transverse scale of 4½/1/2 vs. 4½/1/1. It has longer head length, 29.3(27.5-31.1) vs. 24.5(22.1-26.9)% SL, wider gape

Table 1: Comparative morphometry of *Rasbora ornatus* sp. novo and *R. rasbora*

	<i>Rasbora ornatus</i> sp. novo				<i>Rasbora rasbora</i> (H.B.)	
	Holotype	Paratypes (N=13)			ZSI F 2107/2, 1087	
	MUMF 3032	MUMF 3033/10, 1210/3				
		Range	Mean	S.D.	Range	Mean
Standard length (mm)	56.0	35.1-77.6			65.5-124.1	
% of Standard length						
Body depth	29.3	21.1-29.3	28.0	2.1	26.0-30.5	28.3
Caudal length	27.5	26.3-32.4	30.2	1.7	18.9-28.9	23.9
Head length	27.5	27.5-31.1	29.3	0.9	22.1-26.9	24.5
Head depth (occiput)	17.7	16.8-19.5	18.4	0.6	14.8-17.6	16.2
Head depth (eye)	15.0	13.2-15.2	14.8	0.7	11.6-13.9	12.8
Head width (nares)	10.5	8.9-10.9	10.2	0.6	7.8-8.7	8.3
Max. head width	14.3	13.2-15.7	14.7	0.7	12.5-12.6	12.6
Mouth Gape width	9.1	7.1-9.1	8.1	0.5	6.0-7.6	6.8
Body width (dorsal)	15.5	12.4-15.5	14.7	1.0	10.1-12.7	11.4
Body width (anal)	10.4	7.4-10.7	10.0	1.1	8.6-8.6	8.6
Length of caudal peduncle	17.5	13.8-18.2	15.9	1.5	17.6-20.2	18.9
Height of caudal peduncle	13.0	11.9-14.8	13.8	0.5	13.3-13.4	13.4
Pre-dorsal length	52.5	52.1-55.6	53.3	1.0	52.7-53.9	53.1
Post-dorsal length	49.1	44.2-49.1	46.8	1.5	45.7-46.6	46.2
Pre-pelvic length	51.9	47.6-52.6	51.2	1.1	48.6-49.0	48.8
Pre-anal length	74.6	70.2-75.7	73.8	1.7	67.5-73.6	70.6
Pre-anus length	73.0	69.1-74.1	72.0	1.6	66.1-71.6	68.9
Dorsal fin base length	11.4	10.6-13.1	12.4	0.8	10.4-11.5	11.0
Dorsal fin height	20.5	20.2-25.4	22.3	1.5	17.1-22.4	19.8
Pectoral fin length	18.8	18.8-22.8	21.4	1.2	21.0-21.7	21.4
Pelvic fin length	16.6	16.6-19.8	18.5	0.8	14.6-19.5	17.1
Anal fin base length	7.0	7.0-12.1	9.7	1.3	9.6-10.2	9.9
Anal fin height	17.0	16.0-20.2	18.7	1.3	8.9-17.1	13.0
% of Head Length						
Snout length	29.9	23.2-29.9	26.4	1.8	26.7-27.0	26.9
Eye diameter	24.0	22.9-31.3	25.9	2.3	27.8-30.7	29.3
Inter-orbital width	37.0	29.8-39.0	35.7	2.5	25.6-37.5	31.6
Inter-narial width	25.3	20.8-27.6	24.6	1.8	13.9-21.0	17.5
Other ratios						
ED*/Snout length	80.4	80.4-134.8	94.9	15.1	104.3-113.5	108.9
EDI/OW*	64.9	60.9-93.9	73.1	10.3	74.2-120.0	97.1
LCPI/HCP*	1.3	0.9-1.5	1.2	0.1	1.5-1.9	1.7

\* ED: Eye diameter; IOW: Inter-orbital width, LCP: Length of caudal peduncle; HCP: Height of caudal peduncle

**Table 2:** Morphometry and distribution of *Rasbora* species closely allied to *R. ornatus* sp. novo

Characters	<i>Rasbora ornatus</i> sp. nov	<i>Rasbora rasbora</i> (Ham.-Buch.)	<i>Rasbora daniconius</i> (Ham.-Buch.)	<i>Rasbora labiosa</i> (Muke-i)	
		F2107/2, 1087	Brittan (1954)		
1 Lateral line pores	11-20	28-31	28-31	29-32	23-31
2 Lateral line scales	26-28	28-31	28-31	32-35	33-35
3 Lateral transverse scales	4%/112	4Y:J1/1	4%/1/1 %	4%/1/2%	4%/1/2%
4 Dark coloured lateral longitudinal band	Snout tip to end of median caudal rays (tip of caudal black)	Opercle to caudal fin base	Opercle to caudal fin base	Snout tip to caudal base	Opercle to caudal fin base
5 Circumpeduncular scales	14	13	12	14 13-	14 14-
6 Pre-dorsal scales	12-13	13	12	15	15
7 Pre-pelvic scales	13	15			
8 Pre-anal scales	20	24			
9 Gill rakers	4-5+11	3+9-10		2-3+9-12	
10 Distribution	India: Chindwin drainage in Manipur	India: Gangetic provinces, Coromandel coast. Bangladesh, Myanmar, Thailand	India: Gangetic valley, Assam, Peninsular region. Myanmar, Siam	Widely distributed in Indian region	India: Deolali, Maharashtra state

width of 8.1(7.1-9.1) vs. 6.8(6.0-7.6)% SL. Gill rakers 4-5+ 11 vs. 3+9-10 (Roberts 1989). Circumpeduncular scale rows 14 vs. 12-13. The lateral stripe in *R. ornatus* sp. nov. runs from tip of snout to median caudal rays, whereas it extends from opercle to base of caudal fin in *R. rasbora*. Three specimens of *Rasbora* (MUMF 1210-1212) from Chatrickong R., Ukhrul district, Manipur, identified as *R. rasbora* by Vishwanath *et al.* (1998) have been found to have incomplete lateral lines and therefore belong to the new species.

*R. ornatus* is distinguished from the description given by Brittan (1954) of *R. labiosa* Mukerji of the Daniconius complex in having 11-20 vs. 23-31 lateral line pores, 26-28 vs. 33-35 in the lateral line series, lateral transverse scales  $4\frac{1}{2}/1\frac{1}{2}$  vs.  $4\frac{1}{2}/1\frac{1}{2}\frac{1}{2}$ ; simple lips vs. greatly hypertrophied lower lip, and fleshy sheath on the first dorsal fin ray absent vs. present. Jayaram (1999) considers the latter two characters as important distinguishing features of *R. labiosa* Mukerji from other species.

It does not belong to the Pauciperforata complex, which show incomplete lateral line because of its larger size, 35.1-77.6 vs. less than 55 mm SL, 14 vs. 12 circumpeduncular scale rows.

The specimen of *Rasbora rasbora* (F 2516/2) in Zoological Survey of India, Kolkata, on close examination has 14 anal fin rays, much more than the usual numbers in other species of the genus (5-6). Its dorsal fin is inserted much backward, i.e., nearer the caudal fin base (predorsal length is 66.4% SL and post-dorsal length is 32.7% SL). Brittan (1954) places fishes with such characters in other genera of Danioninae.

#### Comparative materials.

1. *Rasbora rasbora* (Hamilton-Buchanan), F. 2107/2 ZSI, Dharikati, 31,12 miles R. Bharati near Lokra (Balipara F. track, Assam). Dr. S.L. Hora; no date.

2. *R. rasbora*, 1087 ZSI, no collection data.

#### ACKNOWLEDGEMENT

The first author is grateful to ICAR-NATP Project on Germplasm Inventory and Gene-Banking of Freshwater Fishes for financial assistance.

#### REFERENCES

- BRITTAN, M.R. (1954): A revision of the Indo-Malayan freshwater fish genus *Rasbora*. *Monogr. Inst. Sci. & Tech., Manila*, 3: 224.
- HORA, S.L. (1921): Fish and fisheries of Manipur with some observations on those of the Naga Hills. *Rec. Indian Mus.* 22(3): 165-214, 4 pls.
- HORA, S.L. & D.D. MUKERJI (1935): Fishes of the Naga hills, Assam. *Rec. Indian Mus.* 37: 381-404.
- JAYARAM, K.C. (1999): The freshwater fishes of the Indian region. Narendra Publ. House, New Delhi. 551 pp.
- ROBERTS, T.R. (1989): The freshwater fishes of Western Borneo (Kalimantan, Barat, Indonesia). *Mem. Calif Acad. Sci.* 14: 1-210.
- VISHWANATH, W., W. MANOJKUMAR, L. KOSYGIN & K. SELIM (1998): Biodiversity of freshwater fishes of Manipur, India. *Italian J. Zool.* 65: 321-324.

• • •